# LaboPass™ IP pro-Taq PCR Premix

Cat. No. CMT6005



#### Lot No.

## **Description**

- LaboPass™ IP pro-Tag PCR Premix is an optimized 2X PCR Master mix containing IP pro-Tag Polymerase, dNTPs, MgCl<sub>2</sub>, reaction buffer, loading dye and stabilizers that is aliquoted into the Thin-Wall 8-strip PCR tube. This premix formulation simplifies PCR setup. The user simply adds template, primers, and DW to start the reaction.
- LaboPass™ IP pro-Tag DNA Polymerase is a modified version of Tag polymerase which improves the reliability and specificity of PCR reaction. The IP pro-Tag Polymerase has proofreading activity and is more thermostable than wild type Tag DNA polymerase, which allows for the amplification of long length up to 20 kb with high accuracy. The amplified products contain a mixture of blunt ends and 3' A-plus ends.

## **Specifications**

IP pro-Taq Polymerase, dNTPs, reaction buffer, Components

loading dye stabilizers

Ready-to-use Type

(Only DNA template and primers are needed)

Reaction volume

(2X PCR Master mix is aliquoted (each 10 μl) into

the PCR tube)

Store at -20℃

# Storage and Stability

LaboPass<sup>™</sup> PCR Premix is stable for 1 year when stored at -20°C. Repeated freezing and thawing of the premix is not recommended.

# **Applications**

- · General PCR for difficult template
- · Long range PCR
- TA-cloning

## **Quality Control**

Each lot of IP pro-Tag polymerase, reaction buffer and dNTPs is tested for contamination such as *E.coli* genomic DNA, nicking, endonuclease and exonuclease.

## **Standard Reaction**

Components	Volumes (µl)
2X IP pro- <i>Taq</i> PCR Premix	10 µl
Forward Primer (10~50 pmoles/µI)	1 µl
Reverse Primer (10~50 pmoles/µl)	1 µl
DNA Template (Variable*)	1~2 µl
Distilled water	6~7 µl
Total reaction volume	20 µl

#### \* Amount of DNA template

10-200 ng - Eukaryotic genomic DNA 1-50 na - Prokaryotic genomic DNA - Purified homogeneous DNA <5 ng (e.g. plasmid, lambda DNA. etc)

- cDNA: 0.5-10% of RT reaction volume

# **General Thermo-Cycler protocol**

Step	Time	Temperature
Initial denaturation	1-5 min	94-95°C
25-35 Cycles: Denaturation Annealing Extension	10-25 sec 10-25 sec 60 sec/1 kb	94-95°C 45-70°C 68-72°C
Final extension	5 min	68-72°C

#### \* Note

- Vortex all solutions and spin down carefully before using
- Dispense on ice and spin down again